

# Relationship between the [O III] $\lambda$ 5007 line and 5GHz radio emission

A. Labiano<sup>1,2,3</sup>

<sup>1</sup> Instituto de Estructura de la Materia, Serrano 121, 28006, Madrid, Spain

<sup>2</sup> Kapteyn Astronomical Institute, Groningen, 9700 AV, The Netherlands

<sup>3</sup> Rochester Institute of Technology, Rochester, NY, 14623, USA

The dates of receipt and acceptance should be inserted later

**Key words** galaxies: active, galaxies: jets, galaxies: interactions, ISM: jets and outflows.

I have compiled observations of [O III]  $\lambda$  5007 line and 5 GHz radio emission for a large sample of GPS, CSS and FR sources. Several properties were studied and compared. The most relevant findings are that the FWHM and the luminosity of the [O III]  $\lambda$  5007 line are correlated with the size of the radio source. I present the data and discuss the correlations, with special focus on jet-host interaction, triggering and enhancing of [O III]  $\lambda$  5007 emission.

© 2005 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

## 1 Introduction

Current models for the evolution of powerful radio galaxies suggest that these sources propagate from the  $\sim 10$  pc to Mpc scales at roughly constant velocity through an ambient medium which declines in density as  $\rho(R) \propto R^{-2}$  while the sources decline in radio luminosity as  $L_{rad} \propto R^{-0.5}$  (e.g., O’Dea 2002, and references therein). In this scenario, GPS would evolve into CSS and these into large -supergalactic sized- sources<sup>1</sup>. Such a scenario is consistent with the observed number densities of powerful radio sources as a function of linear size (e.g. O’Dea & Baum 1997, Fanti et al. 2001). However, to match observations, the radio jets of the young sources must slow as they cross the host ISM and dim faster than predicted. The most likely mechanism to produce these effects is interaction of the radio source with the host environment.

Interaction was found (e.g., Labiano et al. 2008; Holt et al. 2006; Axon et al. 2000). However, the studies focused in small samples or even just a few sources. Until now, there was no study of interaction of a large representative sample of GPS and CSS sources. Furthermore, the sparseness of samples studied did not allow a general study of the consequences of interaction on the models or the host.

In order to study the interaction between the gas clouds and the radio source I collected a sample of almost one hundred sources, including GPS and CSS galaxies and quasars, as well as FR sources. I study the properties of the [O III]  $\lambda$  5007 line and 5 GHz radio emission looking for traces of interaction and the mechanisms responsible for it.

## 2 The sample

I have compiled an representative sample of GPS, CSS and large sources with observations of the [O III]  $\lambda$  5007 line and 5 GHz radio emission. The sample consists of literature data (O’Dea 1998; Gelderman & Whittle 1994, de Vries et al. 2000), the 2-Jy radio sources (Morganti et al. 1997; di Serego-Alighieri et al. 1994; Morganti et al. 1993; Tadhunter et al. 1993), STIS data (O’Dea et al. 2002; Labiano et al. 2005) and new Kitt Peak observations of three GPS radio galaxies: 0554-026, 0941-080, 1345+125. The total number of sources included in the sample is 95 (21 GPS, 22 CSS, 52 large sources).

## 3 Results and discussion


The following properties of the [O III]  $\lambda$  5007 line and 5 GHz radio emission were compared: FWHM, luminosity, asymmetry, kurtosis ([O III]  $\lambda$  5007 line), power, size and turnover frequency (radio emission) in the sample.

The [O III]  $\lambda$  5007 FWHM shows no correlation with radio power, turnover frequency and [O III]  $\lambda$  5007 luminosity. Therefore, the shock velocity (closely related to the FWHM, see e.g., Bicknell et al. 1997) is independent of the strength radio source and does not affect the luminosity of the ionized gas or the radio spectral properties of the source. However, the data suggests a possible correlation between [O III]  $\lambda$  5007 FWHM and radio source size (Figure 1) suggesting a possible deceleration of the jet as it crosses the host:


$$\log \text{FWHM} \simeq 2.89(\pm 0.04) - 0.08(\pm 0.05) \times \log \text{LS}$$

The correlation is not very strong. However, the deceleration of the jet is required to predict observations (e.g., O’Dea 1998) and the latest models are predicting decelerations in the jet (see e.g., Kawakatu et al. 2009, in this

<sup>1</sup> Recent work added the High Frequency Peakers to the sequence, as possible progenitors of GPS (e.g., Orienti 2007 et al. and references therein).



**Fig. 1** Plot of the [O III]  $\lambda$  5007 FWHM of the sources in Gelderman & Whittle 1994 (diamonds), de Vries et al. 2000 (triangles) our Kitt Peak (circles), STIS (shaded diamonds) observations, and a linear fit to the data. The ionization and kinematics of 3C 67, 3C 277.1 and 3C 303.1 are studied in Labiano et al. (2005) and O’Dea et al. (2002).




**Fig. 2** Radio power at 5GHz versus linear size for GPS, CSS and large radio sources.

volume). Unfortunately, there is not much data available of [O III]  $\lambda$  5007 FWHM.

As expected (e.g., O’Dea & Baum 1997, the sample also shows no correlation between radio power and radio size (Figure 2), showing that GPS, CSS and FR2 sources are equally powerful ( $\log \text{Power}_{5\text{GHz}} \sim 10^{26-27}$ ) while FR1 sources tend to be fainter.

It is also clear that quasars tend to be brighter in [O III]  $\lambda$  5007 radio than galaxies, consistent with Unification scenarios: some of the [O III]  $\lambda$  5007 may be hidden by the torus in radio galaxies (e.g. Hes et al. 1993). However, the difference in [O III]  $\lambda$  5007 emission could be due to selection effects: quasars are usually found at higher redshifts. The sample could be missing fainter quasars with luminosities similar to radio galaxies.

The sample shows the known relation (e.g., Rawlings & Saunders 1991; Baum & Heckman 1989) between [O III]  $\lambda$  5007 luminosity and radio power (Figures 3 and 4) for powerful ( $\log \text{Power}_{5\text{GHz}} > 10^{25}$ ) radio sources. This relation is usually explained as the AGN powering both the ionized



**Fig. 3** Plot of the [O III]  $\lambda$  5007 luminosity versus radio power at 5GHz. Data for the CSS and GPS sources from Gelderman & Whittle (1994); O’Dea (1998), de Vries et al. (2000), the 2-Jy sample and our Kitt Peak observations. Data for the large radio sources from the 2-Jy sample. The IDs of the sources have been updated with Zirbel & Baum (1995). I use “C/G” to name those sources with no clear ID as CSS or GPS.

gas and radio emission. For our sample, the different correlations are:

GPS:

$$\log L_{[\text{O III}]} = 32(\pm 4) + 0.4(\pm 0.1) \times \log P_{5\text{GHz}}$$

CSS:

$$\log L_{[\text{O III}]} = 22(\pm 4) + 0.8(\pm 0.2) \times \log P_{5\text{GHz}}$$

GPS + CSS:

$$\log L_{[\text{O III}]} = 25(\pm 3) + 0.7(\pm 0.1) \times \log P_{5\text{GHz}}$$

GPS + CSS + large:

$$\log L_{[\text{O III}]} = 12(\pm 2) + 1.13(\pm 0.07) \times \log P_{5\text{GHz}}$$

It should be noted that the correlation is not present in fainter radio sources (SDSS, Best 2008, in this volume), suggesting that a different mechanism may be producing the radio and [O III]  $\lambda$  5007 emission in these sources.

There is no evident relation between kurtosis or asymmetry of the [O III]  $\lambda$  5007 line with other properties of the line. Most of the sources have kurtosis lower than 1, suggesting the presence of broad wings in the gas. Whittle (1985) also finds the same for his sample of Seyfert galaxies. Most sources shows asymmetry values close to 0 in their [O III]  $\lambda$  5007 profile, suggesting the cocoon widens or narrows in a symmetric way or there are no major variations between both sides. However, the ground spectra may lack of enough resolution to separate more complex structures that could change the profile of the line.

O’Dea (1998) discovered that in the Gelderman & Whittle (1994) sample, GPS galaxies tend to have lower [O III]  $\lambda$  5007 luminosity than CSS galaxies. However, it was not clear if the trend would be followed by a larger sample with different selection criteria, what was the behaviour for quasars and large FR sources, or the possible consequences

for general radio source evolution<sup>2</sup>. To assess these issues, I tested the trend with the current sample, which also includes GPS and CSS quasars and supergalactic-sized sources. I find that the GPS and CSS sources (galaxies and quasars) show a strong correlation between [O III]  $\lambda$  5007 line luminosity and size of the radio source<sup>3</sup>:

GPS + CSS:

$$\log L_{[\text{O III}]} = 42.43(\pm 0.09) + 0.46(\pm 0.09) \times \log LS_{5\text{GHz}}$$

GPS\* + CSS:

$$\log L_{[\text{O III}]} = 42.44(\pm 0.08) + 0.4(\pm 0.1) \times \log LS_{5\text{GHz}}$$

GPS\* means the complete sample of GPS except the smallest source.

In principle, this correlation could be due to the AGN enhancing both the radio and emission gas luminosities by photoionization. However, for the same radio power, small sources (GPS) are systematically fainter in [O III]  $\lambda$  5007 than larger (CSS) sources.

I propose a scenario where the expansion of the radio source through the host ISM is triggering and/or enhancing the [O III]  $\lambda$  5007 line emission through direct interaction. Some contribution from the AGN light must be present, however, AGN light alone, would not produce a correlation with size. Furthermore, the fact that the correlation disappears for supergalactic-sized sources supports this model: once the radio lobes leave the host, the [O III]  $\lambda$  5007 luminosity drops (Figure 5).

This scenario is also supported with previous observations finding evidence of strong interaction between the jet and surrounding ISM, as well as proof of shock-ionized [O III]  $\lambda$  5007 (e.g., Labiano et al. 2005; O'Dea et al. 2003). Jet-ISM interaction is also found through H I studies (e.g., Labiano et al. 2006; Holt et al. 2006, and references therein) and predicted by jet expansion models (e.g., Jeyakumar et al. 2005; Saxton et al. 2005).


Another interesting enhancing mechanism to consider is that the jet could enhance [O III]  $\lambda$  5007, at least partly, through indirect mechanisms such as jet-induced star formation (Labiano et al. 2008, in prep.). However, new data are needed to study the possible contribution of recently formed stars. The hosts of GPS and CSS sources are usually elliptical galaxies so it is unlikely that the average/normal stellar population of the host has a strong to the [O III]  $\lambda$  5007 emission. These "normal" stars would however not create a correlation with radio jet size (and jet-induced stars would).

The correlation divided by source type is:


GPS:

<sup>2</sup> A direct consequence is that it is more difficult to find optical counterparts of GPS than CSS.

<sup>3</sup> Note that the trend is also suggested by Figure 4 and the different correlations between radio power and [O III]  $\lambda$  5007 luminosity for GPS and CSS sources.



**Fig. 4** As Figure 3, showing only the GPS and CSS sources.



**Fig. 5** Plot of the [O III] luminosity versus linear size of the radio source, showing the correlation for GPS and CSS sources. Data from the same references as figure 3.

$$\log L_{[\text{O III}]} = 42.8(\pm 0.2) + 0.8(\pm 0.2) \times \log LS_{5\text{GHz}}$$

GPS\*:

$$\log L_{[\text{O III}]} = 42.5(\pm 0.4) + 0.5(\pm 0.4) \times \log LS_{5\text{GHz}}$$

CSS:

$$\log L_{[\text{O III}]} = 42.6(\pm 0.2) + 0.2(\pm 0.3) \times \log LS_{5\text{GHz}}$$

The correlation tends to disappear when the sample is divided in different types of sources. This could be due to low statistics or to the smaller range of sizes covered by each type. GPS could show lower [O III]  $\lambda$  5007 due to high obscuration. However, it is more likely that young compact sources are too small to strongly affect their environment (this effect has also been observed in star formation histories of GPS hosts, Labiano et al. 2008). The UV luminosities of GPS seem to be as high as the CSS luminosities. Furthermore, the UV luminosity of GPS sources could be correlated with their radio power (Labiano et al. 2008). These two effects suggest that obscuration is not too strong in GPS sources or, at least, similar to CSS.

Concerning the overall scenario of radio source evolution, where GPS and CSS evolve into the large, supergalactic sized, sources, the visual inspection of Figure 5 suggests that CSS would evolve into FR2. Some authors also found

a possible decreasing trend linking FR2 to FR1 (Best 2008, private communication) with increasing size. However, our sample needs more supergalactic-sized sources to address evolution beyond  $\sim 15 - 20$  kpc. Extensive discussions on the FR2 - FR1 connection can be found on the literature: see e.g., Müller et al. 2004; Best et al. 2005; Wold et al. 2007 and references therein, or classical papers such as Baum et al. (1995) and Zirbel & Baum (1995).

#### 4 Summary of main results and future work

The aim of the project was to deepen our understanding of radio jet-host interaction in young AGN, studying the [O III]  $\lambda$  5007 line and 5 GHz radio emission properties of GPS and CSS sources. I have compiled a large ( $\sim 100$  sources), representative sample of GPS and CSS quasars, combined with FR 1 and FR 2, large sources (to help establish a time evolution line) from published data, as well as new data for three GPS sources.

The main result of the study is that the [O III]  $\lambda$  5007 emission is clearly enhanced by the jet expansion through the host ISM. This is consistent with previous observations as well as numerical models of jet expansion. However, further work is required (Labiano et al. 2008, in prep.):

- The supergalactic-size sample needs to be widened, to establish evolution when the radio lobes leave the host.
- Most likely, the [O III]  $\lambda$  5007 emission is enhanced by the jet through shocks. However, it could also be enhanced by star formation induced by gas compressed by the shock, or a combination of both effects.
- Study of star formation and AGN tracers (such as X-rays, 24 and 70  $\mu$ m dust, line ratios and PAH, etc) to separate and evaluate the different mechanisms and contributions to the [O III]  $\lambda$  5007 emission.
- Apply and improve jet expansion models to reproduce the results.

A parallel result to the [O III]  $\lambda$  5007 - radio size correlation is that it will be more difficult to find optical counterparts of GPS than CSS.

The data suggests a possible deceleration in the jet as it crosses the host ISM, which is required by most radio source evolution models. However, the correlation is too weak and a much wider sample, with higher resolution spectra, is needed.

The sample also reflects some already known results such as quasars showing lower [O III]  $\lambda$  5007 luminosity than radio galaxies, radio power correlated with [O III]  $\lambda$  5007 luminosity, and radio power independent of the size of the radio source.

*Acknowledgements.* This research has made use of the NASA / IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. I would like to thank C.P. O'Dea and P. D. Barthel for many fruitful comments and scientific discussions.

#### References

- Axon, D. J., et al. 2000, *AJ*, 120, 2284  
 Baum, S. A. & Heckman, T. 1989, *ApJ*, 336, 702  
 Baum, S. A., Zirbel, E. L., & O'Dea, C. P. 1995, *ApJ*, 451, 88  
 Best, P. N., et al. 2005, *MNRAS*, 362, 25  
 Bicknell, G. V., Dopita, M. A., & O'Dea, C. P. 1997, *ApJ*, 485, 112  
 Carvalho, J. C. 1994, *A&A*, 292, 392  
 Carvalho, J. C. 1998, *A&A*, 329, 845  
 de Vries, W. H., O'Dea, C. P., Barthel, P. D., & Thompson, D. J. 2000, *A&AS*, 143, 181  
 de Vries, W. H., O'Dea, C. P., Baum, S. A., & Barthel, P. D. 1999, *ApJ*, 526, 27  
 de Vries, W. H., et al. 1997, *ApJS*, 110, 191  
 De Young, D. S. 1993, *ApJ*, 402, 95  
 di Serego-Alighieri, S., Danziger, I. J., Morganti, R., & Tadhunter, C. N. 1994, *MNRAS*, 269, 998  
 Fanti, C., et al. 2001, *A&A*, 369, 380  
 Gelderman, R. & Whittle, M. 1994, *ApJS*, 91, 491  
 Heckman, T. M., Miley, G. K., van Breugel, W. J. M., & Butcher, H. R. 1981, *ApJ*, 247, 403  
 Hes, R., Barthel, P. D., & Fosbury, R. A. E., 1993, *Nature*, 362, 326  
 Holt, J., Tadhunter, C. N., & Morganti, R. 2006, *Astronomische Nachrichten*, 327, 147  
 Jeyakumar, S., Wiita, P. J., Saikia, D. J., & Hooda, J. S. 2005, *A&A*, 432, 823  
 Kawakatu, N. & Kino, M. 2008 in prep.,  
 Labiano, A., O'Dea, C. P., Barthel, P. D., de Vries, W. H., & Baum, S. A. 2008, *A&A*, 477, 491  
 Labiano, A., et al. 2005, *A&A*, 436, 493  
 Labiano, A., et al. 2006, *A&A*, 447, 481  
 Marecki, A., Thomasson, P., Mack, K.-H., & Kunert-Bajraszewska, M. 2006, *A&A*, 448, 479  
 Massey, P., Strobel, K., Barnes, J. V., & Anderson, E. 1988, *ApJ*, 328, 315  
 Morganti, R., Killeen, N. E. B., & Tadhunter, C. N. 1993, *MNRAS*, 263, 1023  
 Morganti, R., Tadhunter, C. N., Dickson, R., & Shaw, M. 1997, *A&A*, 326, 130  
 Müller, S. A. H., et al. 2004, *A&A*, 426, L29  
 O'Dea, C. P. 1998, *PASP*, 110, 493  
 O'Dea, C. P. & Baum, S. A. 1997, *AJ*, 113, 148  
 O'Dea, C. P., de Vries, W. H., Koekemoer, A. M., et al. 2003, *Publications of the Astronomical Society of Australia*, 20, 88  
 O'Dea, C. P., de Vries, W. H., Koekemoer, A. M., et al. 2002, *AJ*, 123, 2333  
 Orienti, M., Dallacasa, D., & Stanghellini, C. 2007, *A&A*, 475, 813  
 Rawlings, S., & Saunders, R. 1991, *Nature*, 349, 138.  
 Saxton, C. J., Bicknell, G. V., Sutherland, R. S., & Midgley, S. 2005, *MNRAS*, 359, 781  
 Spergel, D. N., et al. 2003, *ApJS*, 148, 175  
 Stanghellini, C., et al. 2005, *A&A*, 443, 891  
 Tadhunter, C. N., Morganti, R., di Serego-Alighieri, S., Fosbury, R. A. E., & Danziger, I. J. 1993, *MNRAS*, 263, 999  
 Whittle, M. 1985, *MNRAS*, 213, 1  
 Wold, M., Lacy, M., & Armus, L. 2007, *A&A*, 470, 531  
 Zirbel, E. L. & Baum, S. A. 1995, *ApJ*, 448, 521